

Z-Observables and the Effective Weak Mixing Angle in the MSSM

A. Dedes[†], K. Tamvakis

Division of Theoretical Physics, Physics Department, University of Ioannina, GR-45 110, Greece

A. B. Lahanas

Physics Department, Nuclear and Particle Physics Section, University of Athens, Athens 157 71, Greece

We make a comparison of the predicted effective weak mixing angle, the Z-on resonance asymmetries and the W-boson mass to the LEP and SLD data at their present status. We find that the predicted MSSM values for the effective weak mixing angle are in agreement with the LEP+SLD average value for a “heavy” SUSY breaking scale while we observe an agreement with SLD data in the case of a “light” SUSY breaking scale. The resulting values for the W-boson mass and for the electron left-right asymmetries are compatible with CDF,UA2,DØ and LEP data respectively. Unexpectedly we find that the supersymmetric QCD contributions to the Z-observables tend to vanish everywhere in the $M_{1/2}$ - M_0 plane. Furthermore, values of $M_{1/2}$ which are greater than 500 GeV are favoured by the MSSM if one considers the current experimental value for the strong coupling.

I. INTRODUCTION

The weak mixing angle is defined as a ratio of the gauge couplings

$$\hat{s}^2(Q) = \frac{\hat{g}'^2(Q)}{\hat{g}^2(Q) + \hat{g}'^2(Q)}, \quad (1)$$

where $\hat{g}$ and $\hat{g}'$ are the $SU(2)$ and $U(1)_Y$ gauge couplings respectively. These couplings are running in the sense that they depend on the energy scale Q , and in supersymmetry they are evaluated in the $\overline{DR}$ scheme. The weak mixing angle $\hat{s}^2$ can be easily predicted from a GUT model, *i.e.* $SU(5)$ or from a string derived model like the flipped $SU(5)$. There are many sources of the determination of the weak mixing angle in terms of experimentally measurable quantities. From muon decay for instance and knowing the most accurate parameters, $M_Z = 91.1867 \pm 0.0020 \text{ GeV}$, $\alpha_{EM} = 1/137.036$ and $G_F = 1.16639(1) \times 10^{-5} \text{ GeV}^{-2}$, we can define the $\hat{s}^2$ at M_Z by the expression

$$\hat{s}^2 \hat{c}^2 = \frac{\pi \alpha_{EM}}{\sqrt{2} M_Z^2 G_F (1 - \Delta\hat{\alpha}) \hat{\rho} (1 - \Delta\hat{r}_W)}, \quad (2)$$

where

$$\hat{\rho}^{-1} = 1 - \Delta\hat{\rho} = 1 - \frac{\Pi_{ZZ}(M_Z^2)}{M_Z^2} + \frac{\Pi_{WW}(M_W^2)}{M_W^2}, \quad (3)$$

$$\Delta\hat{r}_W = \frac{\Pi_{WW}(0) - \Pi_{WW}(M_W^2)}{M_W^2} + \hat{\delta}_{VB}, \quad (4)$$

$$\hat{\alpha} = \frac{\alpha_{EM}}{1 - \Delta\hat{\alpha}}. \quad (5)$$

Π 's are the loop contributions to the transverse Z and W gauge bosons self energies and they contain both logarithmic and finite parts. The quantity $\hat{\delta}_{VB}$ contains vertex+box corrections and the term $\Delta\hat{\alpha}$ contributions from light quarks and leptons as well contributions from

heavy particle thresholds *i.e.* top quark, Higgs bosons, superpartner masses. The contributions (3),(4),(5) include logarithmic corrections of the form

$$\log\left(\frac{M_{1/2}^2}{M_Z^2}\right),$$

and if the soft gaugino mass $M_{1/2}$ takes on large values we refer to this quantity as a “large log” quantity. We have proven in [1] that the weak mixing angle behaves like

$$\hat{s}^2 \sim \log\left(\frac{M_{1/2}^2}{M_Z^2}\right). \quad (6)$$

This is clear from Figure 1, where if we increase the value of $M_{1/2}$ then $\hat{s}^2$ takes on large values which means that large logs have not been decoupled from its value. Through this note we always assume the unification of gauge couplings up to two loops and the constraint from the radiative symmetry breaking [2] where the full one loop minimization conditions have been taken into account.




FIG. 1. The extracted value of $\hat{s}^2$ as a function of $M_{1/2}$. The logarithmic behavior resulting from the soft gaugino masses is obvious.

TABLE I. Partial and total contributions to Δk_f , ($f = \text{lepton}, \text{charm}, \text{bottom}$), for two sets of inputs shown at the top. Also shown are the predictions for the effective weak mixing angles and the asymmetries. In the first five rows we display the universal contributions to $10^3 \times \Delta k$ of squarks ($\tilde{q}$), sleptons ($\tilde{l}$), Neutralinos and Charginos ($\tilde{Z}, \tilde{C}$), ordinary fermions and Higgses (The number shown in the middle below the "charm" column refers to "lepton" and "bottom" as well). In the next five rows we display the contributions of gauge bosons as well as the supersymmetric *Electroweak* (EW) and *SQCD* vertex and external fermion wave function renormalization corrections to $10^3 \times \Delta k$.

	$M_0 = 100$ $m_t = 175$	$M_{1/2} = 100$ $\tan b = 4$	$A_0 = 100$ $\mu > 0$	$M_0 = 600$ $m_t = 175$	$M_{1/2} = 600$ $\tan b = 4$	$A_0 = 600$ $\mu > 0$
	<i>lepton</i>	<i>charm</i>	<i>bottom</i>	<i>lepton</i>	<i>charm</i>	<i>bottom</i>
$\tilde{q}$		-4.3481			-10.0677	
$\tilde{l}$		-0.2166			-0.4042	
$\tilde{Z}, \tilde{C}$		7.087			-12.2366	
<i>Fermions</i>		4.8177			4.4290	
<i>Higgs</i>		-0.4390			-1.1523	
<i>Gauge</i>	-3.2491	-3.7027	2.2999	-3.0766	-3.5184	2.2771
<i>Vertex(EW)</i>	-1.9399	2.1535	6.9837	2.8229	5.6099	21.2526
<i>Wave(EW)</i>	-0.1487	-2.210	-8.3524	-2.8280	-5.6037	-21.0202
<i>Vertex(SQCD)</i>	-	0.1217	-1.0491	-	0.2416	-1.0202
<i>Wave(SQCD)</i>	-	-0.1204	1.0057	-	-0.2416	1.0190
$\Delta k (\times 10^2)$	0.1557	0.3137	0.7782	-2.2513	-2.2944	-1.6424
$\sin^2 \theta_f$	0.23019	0.23055	0.23162	0.23145	0.23135	0.23277
$\mathcal{A}_{LR}^f$	0.1575	0.6708	0.9355	0.1476	0.6681	0.9347
$\mathcal{A}_{FB}^f$	0.0186	0.0792	0.111	0.0163	0.0740	0.1035

Although the weak mixing angle $\hat{s}^2$ provides a convenient means to test unification in unified extensions of the Standard Model (SM) or the MSSM it is *not* an experimental quantity. There are several studies in the literature evaluating the weak mixing angle both in the SM [3–7] and in the MSSM [8].

II. THE EFFECTIVE WEAK MIXING ANGLE

LEP collaborations employ the effective weak mixing angle s_f^2 first introduced by Degrandi and Sirlin [3]. Electroweak corrections to the $Zf\bar{f}$ vertex yield the effective Lagrangian which defines the effective weak mixing angle as

$$s_f^2 \equiv \hat{s}^2 \hat{k}_f = \hat{s}^2 (1 + \Delta \hat{k}_f), \quad (7)$$

or in analogous way as we define the eq.(2)

$$s_f^2 c_f^2 = \frac{\pi \alpha_{EM} (1 + \Delta \hat{k}_f) (1 - \frac{\hat{s}^2}{\hat{c}^2} \Delta \hat{k}_f)}{\sqrt{2} M_Z^2 G_F (1 - \Delta \hat{\alpha}) \hat{\rho} (1 - \Delta \hat{r}_W)}, \quad (8)$$

where

$$\Delta \hat{k}_f = \frac{\hat{c}}{\hat{s}} \frac{\Pi_{Z\gamma}(M_Z^2) - \Pi_{Z\gamma}(0)}{M_Z^2} + \delta k_f^{SU\!SY} + \dots \quad (9)$$

$\Pi_{Z\gamma}$ denotes the Z- γ propagator corrections and the term $\delta k_f^{SU\!SY}$ the non-universal corrections resulting from the vertex and wave function renormalization corrections. The non-universal corrections $\delta k_f^{SU\!SY}$ contain both diagrams with electroweak and supersymmetric QCD corrections. There are also other SM contributions which although included in our numerical analysis, they are irrelevant to the discussion here.

The effective weak mixing angle is an experimental measured quantity and its average LEP+SLD value is 0.23152 ± 0.00023 [9]. The difference between the two angles in the SM is

$$\text{SM} : s_l^2 - \hat{s}^2 \lesssim O(10^{-4}),$$

which is less than the error quoted by the experimental groups. This fact is due to the small total contribution from fermions and bosons occurring at the one loop level, in the $\overline{MS}$ scheme and the only dominant contributions are obtained from the two-loop heavy top contributions and three-loop QCD effects.

In the MSSM this picture is changed dramatically. The difference between the two angles can be at one loop,

$$\text{MSSM} : s_l^2 - \hat{s}^2 \lesssim O(10^{-3}).$$

So, in the case of the MSSM the two angles may have completely different values and the difference between them can be much greater than the experimental error. As we have shown in ref. [1] although $\hat{s}^2$ suffers from large logs in the MSSM, this is not the case for the s_f^2 .

So far the situation is clear : GUT analysis is able to predict the weak mixing angle $\hat{s}^2$ and then eq.(7) can translate the result into the experimentally measured quantity, the effective weak mixing angle s_f^2 .

As we have previously mentioned, among the interesting features of the effective weak mixing angle is that large logs get decoupled from its value. In ref. [1] we have shown this fact both analytically and numerically. Here we present only the results of the analytical proof. Math formulae can be found in ref. [1].

A. Analytical Proof of the Decoupling

There are three sources of large logs which affect the value of the weak mixing angle s_f^2 :

1. Gauge boson self energies which feed large logs to the quantities $\Delta\hat{r}_W$, $\hat{\rho}$ and $\Delta\hat{k}_f$. These corrections are canceled against large logs stemming from the electromagnetic coupling $\hat{\alpha}(M_Z)$.
2. Vertex, external wave function renormalizations and box corrections to muon decay which affect $\Delta\hat{r}_W$ through δ_{VB}^{SUSY} . These corrections are cancelled against each other out.
3. Non-universal vertex and external fermion corrections to $Z\bar{f}f$ coupling which affects $\Delta\hat{k}_f$. These corrections are also cancelled against each other out.

We conclude that s_f^2 behaves like $\sim O(\frac{M_Z}{M_{1/2}})$ which is the SUSY contribution from the finite parts. It seems that SUSY follows the well known Appelquist-Carazzone decoupling theorem [10].

B. Numerical Proof of the Decoupling

In order to prove the decoupling of large logs, we have included the full supersymmetric one loop corrections for the universal and non-universal part of eq.(8). In all figures that follows we assume the constraint from the radiative EW symmetry breaking [2] and the universality for soft breaking masses at the unification scale M_{GUT} . In Figure 2 we plot the leptonic effective weak mixing angle s_l^2 as a function of the soft gaugino mass parameter $M_{1/2}$ for a low ($M_0 = 100$ GeV) and high ($M_0 = 600$ GeV) value of the soft scalar parameter, M_0 .




FIG. 2. The effective weak mixing angle s_l^2 versus $M_{1/2}$ for two characteristic values of M_0 . The top quark mass is fixed at $m_t = 175$ GeV.

We obtain that increasing the value of $M_{1/2}$, s_l^2 takes on constant values in the limit $M_{1/2} \rightarrow 900$ GeV independently on the soft squark mass M_0 . Finite corrections of the form $O(\frac{M_Z}{M_{1/2}})$ are significant in the region $M_{1/2} \sim M_0 \lesssim 200$ GeV with the tendency to decrease (increase) the s_l^2 when $M_{1/2}$ is greater(smaller) than M_Z . In addition lowering M_0 we obtain smaller values for the s_l^2 . It is worth noting that MSSM seems to agree with the average LEP+SLD value while in the vicinity of $M_0 \sim M_{1/2} \simeq 100$ GeV, s_l^2 agrees with the SLD experimental value*. The parameters A_0 and $\tan\beta^\dagger$ do not affect significantly the extracted value of s_l^2 .

Going deeper, we display in Table I the partial and total contributions to Δk_f . Also shown are the predictions for the effective weak mixing angle s_f^2 and the asymmetries.

We would like to stress five interesting points on this Table :

- The bulk of the SUSY corrections to $\Delta\hat{k}_f$ is carried by the universal corrections which are sizeable due to their dependence on large log terms.
- The contribution of Higgses which is small mimics that of the SM with a mass in the vicinity of ~ 100 GeV.

*The LEP average $s_{eff}^2 = 0.23199 \pm 0.00028$ differs by 2.9σ from SLD value $s_{eff}^2 = 0.23055 \pm 0.00041$ obtained from the single measurement of left-right asymmetry.

[†]The independent parameter $\tan\beta$ plays an important role in the case of s_b^2 .

- Gauge[‡] and Higgs boson contributions tend to cancel large universal contributions of matter fermions.
- We find that the non-universal Electroweak SUSY corrections are very small for a “heavy” SUSY breaking scale, $M_{1/2} = M_0 = A_0 = 600$ GeV. Although separately vertex and external fermion corrections are large they cancel each other out yielding contributions almost two orders of magnitude smaller than the rest of the EW corrections. For a “light” SUSY breaking scale, $M_{1/2} = M_0 = A_0 = 100$ GeV this cancellation does not occur due to the large splittings of the squarks and sleptons.
- The non-universal SQCD contributions, although a priori expected to be larger than the electroweak corrections turn out to be even smaller. This fact is due to cancellations both from its diagram in vertex or in wave function renormalization, and from different types of the diagrams. This situation is valid both in the “heavy” and “light” SUSY breaking limit and we believe that this cancellations arise from a QED like Ward identities between vertex and wave function renormalization corrections.

III. Z-BOSON OBSERVABLES IN THE MSSM

Having obtained the value of the effective weak mixing angle, we are able to find other Z-observables in the MSSM. For instance the physical W-boson mass is defined by

$$M_W = M_Z \hat{c} \sqrt{\hat{\rho}} = M_Z c_f \sqrt{\frac{\hat{\rho}}{1 - \frac{\hat{s}^2}{\hat{c}^2} \Delta \hat{k}_f}}. \quad (10)$$

In figure 3 we plot the W-boson mass versus the input values for the $M_{1/2}$ parameters for two characteristic values of M_0 in the “light” SUSY breaking scale ($M_0 = 100$ GeV) and in the “heavy” SUSY breaking scale ($M_0 = 600$ GeV). As one can see the W-mass is in agreement with the presently experimentally observed value, $M_W = 80.427 \pm 0.075$ ($M_W = 80.405 \pm 0.089$) GeV obtained from LEP(CDF,UA2,DØ) experiments [14] for rather low(high) values of $M_{1/2}$. Moreover, our results are in agreement with those of refs. [11–13].

[‡]Gauge boson contribution are different for the different fermion species l, c, b . This is due to the fact that their non-universal corrections depend on the charge and weak isospin assignments of the external fermions. In the case of the bottom quark there are additional, significant top quark corrections.




FIG. 3. Predicted values for the W-boson mass in the MSSM.

Having at hand the effective weak mixing angle we can easily derive the left-right asymmetries which in term of the s_f^2 can be defined as

$$A_{LR}^f = \mathcal{A}^f = \frac{2 T_3^f (T_3^f - 2 s_f^2 Q^f)}{T_3^{f2} + (T_3^f - 2 s_f^2 Q^f)^2}, \quad (11)$$

where Q^f is the electric charge and T_3^f is the third component of the isospin of the fermion f . As it is depicted in Figure 4, the MSSM predictions for $\mathcal{A}_e$ agree with LEP+SLD average value ($\mathcal{A}_e = 0.1505 \pm 0.0023$) when both $M_{1/2}$ and M_0 take on values around M_Z . In the heavy SUSY breaking scale, the MSSM agrees with the LEP value, $\mathcal{A}_e = 0.1461 \pm 0.0033$. Note that increasing the value of $M_{1/2}$, $\mathcal{A}_e$ takes on constant values which means that large logs have been decoupled from the expression (11). In this case we have recovered SM results displayed in Particle Data Booklet [14].




FIG. 4. Predicted values for the electron left right asymmetry in the MSSM.

So far we have not considered the constraint resulting from the experimental value of $\alpha_s(M_Z)$ [15]. In Figure 5 we have plotted the acceptable values of the soft breaking parameters which are compatible with LEP ($\alpha_s(M_Z) = 0.119 \pm 0.004$, $s_{eff}^2 = 0.23152 \pm 0.00023$) [9] and the CDF/DØ ($m_t = 175 \pm 5$ GeV) data [16].




FIG. 5. Acceptable values in the $M_{1/2} - M_0$ plane according to the LEP+SLD data. Only large values of $M_{1/2}$ are acceptable.

The trilinear soft couplings as well as the parameter $\tan\beta(M_Z)$ are taken arbitrarily in the region (0-900 GeV) and (2-30), respectively. As we observe from Figure 5, MSSM with unification of gauge coupling constants, universality of the soft masses at M_{GUT} and radiative EW symmetry breaking is valid in the region $M_{1/2} \gtrsim 500$ GeV and $M_0 \gtrsim 70$ GeV. The lower bound on $M_{1/2}$ is put mainly from the experimental value of α_s and on M_0

from the requirement that LSP is neutral. In this region, the physical gluino mass is above 1 TeV, the LSP is $\gtrsim 200$ GeV, the squark and slepton masses are $\gtrsim 800$ GeV and $\gtrsim 210$ GeV and the light Higgs boson mass is greater than 108 GeV.

IV. CONCLUSIONS

In ref. [1] we have included the full one-loop supersymmetric corrections to the effective weak mixing angle which is experimentally determined in LEP and SLD experiments. Our analysis enables one to pass from the “theoretical” weak mixing angle $\hat{s}^2$, which can be predicted from GUT analysis, to the experimental effective weak mixing angle s_f^2 . We conclude that :

- There are no logarithmic corrections of the form $\log(\frac{M_{1/2}^2}{M_Z^2})$ (or more generally $\log(\frac{M_{SUSY}^2}{M_Z^2})$) to the effective weak mixing angle.
- Supersymmetric QCD corrections tend to vanish everywhere in the $M_{1/2} - M_0$ plane.
- The predicted MSSM values for the effective weak mixing angle are in agreement with the present LEP+SLD average value in the “heavy” SUSY breaking scale while they are in agreement with the SLD data in the “light” SUSY breaking scale..
- MSSM predicts values of the W-boson mass which are in agreement with the new CDF,UA2,DØ average value.
- In the “heavy” SUSY breaking limit MSSM seems to prefer the experimental LEP value for the electron left-right asymmetry $\mathcal{A}_e$.
- Finally, values of $M_{1/2}$ which are greater than 500 GeV are favoured by the MSSM if one assumes the present LEP+SLD and CDF/DØ data for s_{eff}^2 , α_s and m_t , respectively.

ACKNOWLEDGMENTS

A.D. and K.T. acknowledge financial support from the research program IIENEΔ-95 of the Greek Ministry of Science and Technology. A.B.L. and K. T. acknowledge support from the TMR network “Beyond the Standard Model”, ERBFMRXCT-960090. A. B. L. acknowledges support from the Human Capital and Mobility program CHRX-CT93-0319.

[†] Talk given at International Workshop of the Hellenic Society for the Study of High-Energy Physics, Athens, Greece, 9-11 Apr 1998.

- [1] A. Dedes, A. B. Lahanas, K. Tamvakis, **hep-ph/9801425**, to appear in *Phys. Rev. D*.
- [2] J. Ellis, J. Hagelin, D.V. Nanopoulos and K. Tamvakis, *Phys. Lett.* **125B**, 1983 (275);
L. E. Ibanez and G. G. Ross, *Phys. Lett.* **110B**, 215 (1982);
L. Alvarez-Gaume, J. Polchinski and M. Wise, *Nucl. Phys.* **B221**, 495 (1983);
- [3] G. Degrassi and A. Sirlin, *Nucl. Phys.* **B352**, 342 (1991).
- [4] P. Gambino and A. Sirlin, *Phys. Rev. D* **49**, 1160 (1994).
- [5] W. J. Marciano and J. L. Rosner, *Phys. Rev. Lett.* **65**, 2963 (1990) and Erratum *ibid.* **68**, 898 (1992);
G. Degrassi, S. Fanchiotti and A. Sirlin, *Nucl. Phys.* **B351**, 49 (1991);
B. A. Kniehl and A. Sirlin, *Phys. Rev. D* **47**, 883 (1993);
S. Fanchiotti, B. A. Kniehl and A. Sirlin, *Phys. Rev. D* **48**, 307 (1993);
B. A. Kniehl, presented at Tennessee International Symposium on Radiative Corrections: Status and Outlook, Gatlinburg, TN, 27 Jun - 1 Jul 1994;
A. Sirlin, *Phys. Lett.* **348B**, 201 (1995), addendum-*ibid.* **B352**, 498 (1995);
B. A. Kniehl and A. Sirlin, *Nucl. Phys.* **B458**, 35 (1996);
P. Gambino, *Acta. Phys. Polon.* **B27**, 3671 (1996);
G. Degrassi, P. Gambino and A. Sirlin, *Phys. Lett.* **394B**, 188 (1997).
- [6] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci and A. Vicere, *Phys. Lett.* **288B**, 95 (1992) ; *Nucl. Phys.* **B409**, 105 (1993);
J. Fleischer, O. V. Tarasov and F. Jegerlehner, *Phys. Lett.* **319B**, 249 (1993);
G. Buchalla and A. J. Buras, *Nucl. Phys.* **B398**, 285 (1993);
L. Andeev et al, *Phys. Lett.* **336B**, 560 (1994), E: *ibid.* **B349** 597 (1995);
K. G. Cheturkin, J. H. Kühn and M. Steinhauser, *Phys. Rev. Lett.* **75**, 3394 (1995);
K. G. Cheturkin, J. H. Kühn and A. Kwiatkowski, in: *Precision Calculations for the Z resonance*, CERN 95-03, eds. D. Bardin, W. Hollik, G. Passarino, S. Peris, A. Santamaria, CERN-TH-95-21 (1995);
G. Degrassi, P. Gambino and A. Vicini, *Phys. Lett.* **383B**, 219 (1996).
- [7] A. Djouadi and C. Verzegnassi, *Phys. Lett.* **195B**, 265 (1987);
A. Djouadi, *Nuovo Cimento* **100A** (357) 1988;
B. A. Kniehl, *Nucl. Phys.* **B347**, 86 (1990);
A. Djouadi and P. Gambino, *Phys. Rev. D* **49**, 3499 and 4705 (1994), *ibid.* **D51** 218 (1995), E: *ibid.* **D53** 4111 (1996).
- [8] M. Drees, K. Hagiwara, and A. Yamada, *Phys. Rev. D* **45**, 1725 (1992);
G. Altarelli, R. Barbieri, and F. Caravaglios, *Phys. Lett.* **314B**, 357 (1993);
P. Langacker and N. Polonski, *Phys. Rev. D* **47**, 4028 (1993); **D52** 3081 (1995);
N. Polonski, Report No. UPR-0641-T, **hep-ph/9411378**;
D. Garcia and J. Sola, *Phys. Lett.* **354B**, 335 (1995);
G.L. Kane, R.G. Stuart and J.D. Wells, *Phys. Lett.* **354B**, 350 (1995);
P. H. Chankowski, Z. Plucienik and S. Pokorski, *Nucl. Phys.* **B439**, 23 (1995);
P.H. Chankowski and S. Pokorski, *Phys. Lett.* **366B**, 188 (1996);
P. H. Chankowski, Z. Plucienik, S. Pokorski and C. E. Vayonakis, *Phys. Lett.* **358B**, 264 (1995);
D. Garcia and J. Solà, *Mod. Phys. Lett.* **A9**, 211 (1994);
D. Garcia, R.A. Jiménez and J. Solà, *Phys. Lett.* **347B**, 309 (1995), *ibid.* **B347** 321 (1995), and Erratum *ibid.* **B351** 602 (1995);
D.M. Pierce and J. Erler, **hep-ph/9708374**, talk presented at the 5th International Conference on Physics Beyond the Standard Model, Balholm (1997);
P. H. Chankowski, J. Ellis and S. Pokorski, **hep-ph/9712234**;
W. de Boer *et al.*, **hep-ph/9712376**;
D.M. Pierce and J. Erler, **hep-ph/9801238**.
- [9] G. Altarelli, R. Barbieri and F. Caravaglios, **hep-ph/9712368**.
- [10] T. Appelquist and J. Carazzone, *Phys. Rev. D* **11**, 2856 (1975).
- [11] J. Bagger, K. Matchev, D. Pierce and R. Zhang, *Nucl. Phys.* **B491**, 3 (1997).
- [12] W. de Boer, A. Dabelstein, W. Hollik, W. Mosle, U. Schwickerath, *Z. Phys.* **C75**, 627 (1997);
W. Hollik in: *Electroweak Precision Observables in the MSSM and Global Analysis of Precision Data*, Talk at the International workshop on Quantum Effects in the MSSM, Barcelona 9-23 September 1997.
- [13] P. Chankowski and S. Pokorski, **hep-ph/9707497**, to appear in "Perspectives in supersymmetry" edited by G.L. Kane, World Scientific, 1997.
- [14] R. M. Barnett *et al.*, *Phys. Rev. D* **54**, 1 (1996), and 1997 off-year partial update for the 1998 edition available on the PDG WWW pages (URL: <http://pdg.lbl.gov/>).
- [15] A. Dedes, A. B. Lahanas, J. Rizos and K. Tamvakis, *Phys. Rev. D* **55**, 2955 (1997);
A. Dedes, *PhD Thesis*, **hep-ph/9803339**.
- [16] DØ Collaboration: S. Abachi *et al.*, *Phys. Rev. Lett.* **79**, 1197 (1997);
CDF Collaboration: F. Abe *et al.*, *Phys. Rev. Lett.* **79**, 1992 (1997).